

Pediatric Symptom Checklist

INSTRUCTIONS FOR SCORING

The Pediatric Symptom Checklist is a psychosocial screen designed to facilitate the recognition of cognitive, emotional, and behavioral problems so that appropriate interventions can be initiated as early as possible. Included here are two versions, the parent-completed version (PSC) and the youth self-report (Y-PSC). The Y-PSC can be administered to adolescents ages 11 and up.

The PSC consists of 35 items that are rated as "Never," "Sometimes," or "Often" present and scored 0, 1, and 2, respectively. The total score is calculated by adding together the score for each of the 35 items. For children and adolescents ages 6 through 16, a cutoff score of 28 or higher indicates psychological impairment. For children ages 4 and 5, the PSC cutoff score is 24 or higher (Little et al., 1994; Pagano et al., 1996). The cutoff score for the Y-PSC is 30 or higher. Items that are left blank are simply ignored (i.e., score equals 0). If four or more items are left blank, the questionnaire is considered invalid.

HOW TO INTERPRET THE PSC OR Y-PSC

A positive score on the PSC or Y-PSC suggests the need for further evaluation by a qualified health (e.g., M.D., R.N.) or mental health (e.g., Ph.D., L.I.C.S.W.) professional. Both false positives and false negatives occur, and only an experienced health professional should interpret a positive PSC or Y-PSC score as anything other than a suggestion that further evaluation may be helpful. Data from past studies using the PSC and Y-PSC indicate that two out of three children and adolescents who screen positive on the PSC or Y-PSC will be correctly identified as having moderate to serious impairment in psychosocial functioning. The one child or adolescent "incorrectly" identified usually has at least mild impairment, although a small percentage of children and adolescents turn out to have very little or no impairment (e.g., an adequately functioning child or adolescent of an overly anxious parent). Data on PSC and Y-PSC negative screens indicate 95 percent accuracy, which, although statistically adequate, still means that 1 out of 20 children and adolescents rated as functioning adequately may actually be impaired. The inevitability of both false-positive and false-negative screens underscores the importance of experienced clinical judgment in interpreting PSC scores. Therefore, it is especially important for parents or other laypeople who administer the form to consult with a licensed professional if their child receives a PSC or Y-PSC positive score.

For more information, visit the Web site: <http://psc.partners.org>.

REFERENCES

- Jellinek MS, Murphy JM, Little M, et al. 1999. Use of the Pediatric Symptom Checklist (PSC) to screen for psychosocial problems in pediatric primary care: A national feasibility study. *Archives of Pediatric and Adolescent Medicine* 153(3):254-260.
- Jellinek MS, Murphy JM, Robinson J, et al. 1988. Pediatric Symptom Checklist: Screening school-age children for psychosocial dysfunction. *Journal of Pediatrics* 112(2):201-209. Web site: <http://psc.partners.org>.
- Little M, Murphy JM, Jellinek MS, et al. 1994. Screening 4- and 5-year-old children for psychosocial dysfunction: A preliminary study with the Pediatric Symptom Checklist. *Journal of Developmental and Behavioral Pediatrics* 15:191-197.
- Pagano M, Murphy JM, Pedersen M, et al. 1996. Screening for psychosocial problems in 4-5 year olds during routine EPSDT examinations: Validity and reliability in a Mexican-American sample. *Clinical Pediatrics* 35(3):139-146.

Pediatric Symptom Checklist (PSC)

Emotional and physical health go together in children. Because parents are often the first to notice a problem with their child's behavior, emotions, or learning, you may help your child get the best care possible by answering these questions. Please indicate which statement best describes your child.

Please mark under the heading that best describes your child:

		Never	Sometimes	Often
1. Complains of aches and pains	1	_____	_____	_____
2. Spends more time alone	2	_____	_____	_____
3. Tires easily, has little energy	3	_____	_____	_____
4. Fidgety, unable to sit still	4	_____	_____	_____
5. Has trouble with teacher	5	_____	_____	_____
6. Less interested in school	6	_____	_____	_____
7. Acts as if driven by a motor	7	_____	_____	_____
8. Daydreams too much	8	_____	_____	_____
9. Distracted easily	9	_____	_____	_____
10. Is afraid of new situations	10	_____	_____	_____
11. Feels sad, unhappy	11	_____	_____	_____
12. Is irritable, angry	12	_____	_____	_____
13. Feels hopeless	13	_____	_____	_____
14. Has trouble concentrating	14	_____	_____	_____
15. Less interested in friends	15	_____	_____	_____
16. Fights with other children	16	_____	_____	_____
17. Absent from school	17	_____	_____	_____
18. School grades dropping	18	_____	_____	_____
19. Is down on him or herself	19	_____	_____	_____
20. Visits the doctor with doctor finding nothing wrong	20	_____	_____	_____
21. Has trouble sleeping	21	_____	_____	_____
22. Worries a lot	22	_____	_____	_____
23. Wants to be with you more than before	23	_____	_____	_____
24. Feels he or she is bad	24	_____	_____	_____
25. Takes unnecessary risks	25	_____	_____	_____
26. Gets hurt frequently	26	_____	_____	_____
27. Seems to be having less fun	27	_____	_____	_____
28. Acts younger than children his or her age	28	_____	_____	_____
29. Does not listen to rules	29	_____	_____	_____
30. Does not show feelings	30	_____	_____	_____
31. Does not understand other people's feelings	31	_____	_____	_____
32. Teases others	32	_____	_____	_____
33. Blames others for his or her troubles	33	_____	_____	_____
34. Takes things that do not belong to him or her	34	_____	_____	_____
35. Refuses to share	35	_____	_____	_____

Total score _____

Does your child have any emotional or behavioral problems for which she or he needs help?

() N () Y

Are there any services that you would like your child to receive for these problems?

() N () Y

If yes, what services? _____

Pediatric Symptom Checklist—Youth Report (Y-PSC)

Please mark under the heading that best fits you:

		Never	Sometimes	Often
1. Complain of aches or pains	1	_____	_____	_____
2. Spend more time alone	2	_____	_____	_____
3. Tire easily, little energy	3	_____	_____	_____
4. Fidgety, unable to sit still	4	_____	_____	_____
5. Have trouble with teacher	5	_____	_____	_____
6. Less interested in school	6	_____	_____	_____
7. Act as if driven by motor	7	_____	_____	_____
8. Daydream too much	8	_____	_____	_____
9. Distract easily	9	_____	_____	_____
10. Are afraid of new situations	10	_____	_____	_____
11. Feel sad, unhappy	11	_____	_____	_____
12. Are irritable, angry	12	_____	_____	_____
13. Feel hopeless	13	_____	_____	_____
14. Have trouble concentrating	14	_____	_____	_____
15. Less interested in friends	15	_____	_____	_____
16. Fight with other children	16	_____	_____	_____
17. Absent from school	17	_____	_____	_____
18. School grades dropping	18	_____	_____	_____
19. Down on yourself	19	_____	_____	_____
20. Visit doctor with doctor finding nothing wrong	20	_____	_____	_____
21. Have trouble sleeping	21	_____	_____	_____
22. Worry a lot	22	_____	_____	_____
23. Want to be with parent more than before	23	_____	_____	_____
24. Feel that you are bad	24	_____	_____	_____
25. Take unnecessary risks	25	_____	_____	_____
26. Get hurt frequently	26	_____	_____	_____
27. Seem to be having less fun	27	_____	_____	_____
28. Act younger than children your age	28	_____	_____	_____
29. Do not listen to rules	29	_____	_____	_____
30. Do not show feelings	30	_____	_____	_____
31. Do not understand other people's feelings	31	_____	_____	_____
32. Tease others	32	_____	_____	_____
33. Blame others for your troubles	33	_____	_____	_____
34. Take things that do not belong to you	34	_____	_____	_____
35. Refuse to share	35	_____	_____	_____